

DYSGLYCEMIA-BASED CHRONIC DISEASE: AN AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS POSITION STATEMENT

*Jeffrey I. Mechanick, MD, FACP, FACE, FACN, ECNU¹; Alan J. Garber, MD, PhD, FACE²;
George Grunberger, MD, FACP, FACE³; Yehuda Handelsman, MD, FACP, FNLA, MACE⁴;
W. Timothy Garvey, MD, FACE⁵*

This document represents the official position of the American Association of Clinical Endocrinologists and the American College of Endocrinology. Where there were no randomized controlled trials or specific U.S. FDA labeling for issues in clinical practice, the participating clinical experts utilized their judgment and experience. Every effort was made to achieve consensus among the committee members. Position statements are meant to provide guidance, but they are not to be considered prescriptive for any individual patient and cannot replace the judgment of a clinician.

Submitted for publication June 28, 2018

Accepted for publication July 31, 2018

From ¹The Marie-Josée and Henry R. Kravis Center for Clinical Cardiovascular Health at Mount Sinai Heart, New York, New York, ²Baylor College of Medicine, Division of Endocrinology, Houston, Texas, ³Wayne State University School of Medicine, Bloomfield Hills, Michigan, ⁴Metabolic Institute of America, Tarzana, California, and ⁵Department of Nutrition Sciences, University of Alabama at Birmingham, Birmingham, Alabama.

Address correspondence to Dr. Jeffrey I. Mechanick, 1 Gustave L. Levy Place, Box 1030, New York, NY 10029.

E-mail: jeffreymechanick@gmail.com.

Published as a Rapid Electronic Article in Press at <http://www.endocrinepractice.org>. DOI: 10.4158/PS-2018-0139

To purchase reprints of this article, please visit: www.aace.com/reprints.

Copyright © 2018 AACE.

This material is protected by US copyright law. To purchase commercial reprints of this article, visit www.aace.com/reprints. For permission to reuse material, please access www.copyright.com or contact the Copyright Clearance Center, Inc. (CCC).

AACE Diabetes Scientific Committee Reviewers

Yehuda Handelsman, MD, FACP, FNLA, FACE, Chair
 David Bell, MB, FACP, FACE
 Lawrence Blonde, MD, FACP, MACE
 Zachary Bloomgarden, MD, MACE
 Stephen Brunton, MD
 Michael Bush, MD
 Samuel Dagogo-Jack, MD, DM, FRCP, FACE
 Jaime Davidson, MD, FACP, MACE
 Ralph DeFronzo, MD, BMS, MS, BS
 Daniel Einhorn, MD, FACP, FACE
 Vivian Fonseca, MD, FACE
 Alan Garber, MD, PhD, FACE
 Ved Gossain, MD, FRCP, MACP, FACE
 George Grunberger, MD, FACP, FACE
 Irl Hirsch, MD
 Paul Jellinger, MD, MACE
 Rosemary Lajara, MD
 Philip Levy, MD, MACE
 Janet McGill, MD, FACE
 John Miles, MD
 Etie Moghissi, MD, FACP, FACE
 Eric Orzech, MD, FACP, FACE
 Leigh Perreault, MD
 S. Sethu Reddy, MD, MBA, FRCPC, FACP, MACE
 Victor Roberts, MD, MBA, FACP, FACE
 Paul Rosenblit, MD, PhD, FNLA, FACE
 Lance Sloan, MD, MS, FASN, FACE
 Guillermo Umpierrez, MD, FACP, FACE
 Jeff Unger, MD, FACE
 Aaron Vinik, MD, PhD, FCP, MACP, FACE
 Farhad Zangeneh, MD, FACP, FACE

ABSTRACT

The American Association of Clinical Endocrinologists (AACE) has created a dysglycemia-based chronic disease (DBCD) multimorbidity care model consisting of four distinct stages along the insulin resistance-prediabetes-type 2 diabetes (T2D) spectrum that are actionable in a preventive care paradigm to reduce the potential impact of T2D, cardiometabolic risk, and cardiovascular events. The controversy of whether there is value, cost-effectiveness, or clinical benefit of diagnosing and/or managing the prediabetes state is resolved by regarding the problem, not in isolation, but as an intermediate stage in the continuum of a progressive chronic disease with opportunities for multiple concurrent prevention strategies. In this context, stage 1 represents “insulin resistance,” stage 2 “prediabetes,” stage 3 “type 2 diabetes,” and stage 4 “vascular complications.” This model encourages earliest intervention focusing on structured lifestyle change. Further scientific research may eventually reclassify stage 2 DBCD prediabetes from a predisease to a true disease state. This position statement is consistent with a portfolio of AACE endocrine disease

care models, including adiposity-based chronic disease, that prioritize patient-centered care, evidence-based medicine, complexity, multimorbid chronic disease, the current health care environment, and a societal mandate for a higher value attributed to good health. Ultimately, transformative changes in diagnostic coding and reimbursement structures for prediabetes and T2D can provide improvements in population-based endocrine health care. (**Endocr Pract.** 2018;24:995-1011)

Abbreviations:

A1C = hemoglobin A1c; **AACE** = American Association of Clinical Endocrinologists; **ABCD** = adiposity-based chronic disease; **CVD** = cardiovascular disease; **DBCD** = dysglycemia-based chronic disease; **FPG** = fasting plasma glucose; **GLP-1** = glucagon-like peptide-1; **MetS** = metabolic syndrome; **T2D** = type 2 diabetes

THE PREDIABETES PROBLEM

The earliest record of diabetes dates back to the Egyptian physician “Hesy-Ra” around 3000 BC (1). Through the years, diabetes has been diagnosed by physical symptoms (e.g., excessive thirst and urination [1]), the presence of overt glycosuria, numerical values representing blood glucose levels over two standard deviations from the mean (2,3), glucose levels based on an epidemiologic association with the development of retinopathy (4), time-averaged measures of dysglycemia for 2 weeks (e.g., glycated protein [fructosamine]) or 2-months (e.g., hemoglobin A1c [A1C]), novel markers and their combinations (5-7), the application of continuous glucose monitoring (CGM) to optimize glycemic control (8), and even genetic variants (9). According to the 2014 National Diabetes Statistics Report by the Centers for Disease Control and Prevention, type 2 diabetes (T2D) accounted for about 90 to 95% of the 29.1 million people of all ages diagnosed and undiagnosed with diabetes in the U.S., with another 86 million people over age 18 years with prediabetes, of which 15 to 30% developed T2D within 5 years (10). However, in the most recent 2017 National Diabetes Statistics Report, these numbers increased and were 30.3 million U.S. residents of all ages (9.4% of the population) with diabetes and 84.1 million age 18 and over (33.9% of the population) with prediabetes (11).

According to a recent analysis using data from the U.S. National Health and Nutrition Examination Surveys (NHANES; 1988-2014), patients with prediabetes have increased prevalence rates of hypertension, dyslipidemia, chronic kidney disease, and cardiovascular disease (CVD) risk (12). Since intervention once a patient develops T2D is more costly, both economically and in terms of disease burden, and lifestyle interventions introduced as early as possibly in the natural history of the disease can be effec-

tive and durable, there is high impact and opportunity associated with a diagnosis of prediabetes (12-18). However, what is particularly alarming is that in 2015, 23.8% of patients with diabetes (7.2 million persons) and 88.4% of patients with prediabetes (74.3 million persons) did not even know they had the condition (11). Generally speaking, this problem of “unawareness” amplifies the impact of chronic disease and emphasizes the importance of identifying specific causes, interventional strategies (including all types of prevention), and implementation tactics. In sum, the critical question is whether making the diagnosis of “prediabetes” can improve health?

Prediabetes was first described in 1956 in the perspective of gestational diabetes (18) but is now a construct in clinical medicine that identifies people at higher risk for T2D, hypertension, and CVD than the general population. Prediabetes is diagnosed based on fasting plasma glucose (FPG), standardized 2-hour postchallenge glucose testing, or A1C levels, though diagnostic cutoffs may vary (19-21). The concept of prediabetes hinges on a preventive care approach to chronic disease, in which people with a “predisease” are identified by a positive screening test, with the explicit intent to intervene. At the population level, the intended result of the intervention would result in decreased incidence and prevalence rates, costs, and detriment on quality-of-life metrics (22). Hence, those with overweight are at higher risk for obesity, prehypertension for hypertension, metabolic syndrome (MetS) for CVD, and so forth.

From a purely theoretical and idealistic standpoint, the advantages of detecting and intervening in people at-risk for a potentially debilitating chronic disease such as T2D makes sense, not only for the individual person but also for society as a whole (22,23). In the 15-year follow-up of the Diabetes Prevention Program Outcomes Study (24), those patients with prediabetes randomized to lifestyle intervention had significant reduction in T2D development. Moreover, among all patients in the study, there was a 28% lower prevalence of microvascular complications in those not developing T2D (24). Unfortunately, from a purely pragmatic standpoint, the actual effects of this type of population-based intervention may prove expensive due to the high prevalence rate of prediabetes, the small number of individuals destined to develop clinically significant T2D, and lack of anticipated level of benefit for all people in the current prediabetes diagnostic category (25). The problem is even deeper. The current paradigm of diabetes care is plagued by patient adherence problems, inertia among health care professionals adopting contemporary clinical practice guidelines, and a lack of reverent, consistent, and universal insurance coverage for diagnostics, lifestyle medicine, and pharmaceuticals, all prompting the need for a new approach involving patient-centered and preventive care (26). In other words, the current management of patients with prediabetes should be challenged,

with diagnostic criteria reworked, logistics of care re-engineered, and the pathophysiologic basis re-examined (27).

The American Association of Clinical Endocrinologists (AACE) has long supported clinical intervention in patients with prediabetes (28-30), together with evidence-based optimization of care for patients with diabetes, obesity, and related metabolic disorders (31-34). With respect to obesity, AACE has recently proposed a new Adiposity-Based Chronic Disease (ABCD) diagnostic term (35) and a complications-centric obesity care model. Now, within a larger framework of comprehensive cardiometabolic risk, this thinking extends to dysglycemia.

AACE recognizes the dilemma confronting prediabetes as an actionable diagnosis for intervention and resolves this difficulty by redefining the context of prediabetes and T2D within an evidentiary structure for *Dysglycemia-Based Chronic Disease (DBCD)*. Rather than strongly arguing that DBCD should immediately be part of our lexicon, AACE takes the position that popular and recognizable terms, such as prediabetes, T2D, and MetS, fit squarely within a DBCD framework as evolving manifestations of a chronic disease. Therefore, research, education, and clinical practice processes should re-orient toward effective prevention and treatment at all stages in this chronic disease model. More to the point, that despite objections and controversy already anchored in the literature (36,37), AACE takes the firm stance that patients with prediabetes fall within the progressive spectrum of DBCD (insulin resistance-prediabetes-T2D) and are well-suited for structured lifestyle and/or pharmaco-therapeutic preventive measures. Notably, this effort to better reconcile prediabetes within a larger framework of cardiometabolic health is shared by others, where the targets are precise, classification is rational, and early intervention improves clinical outcomes (38,39).

THE GENERAL CONTEXT OF DBCD WITHIN THE NEXUS OF ABCD AND CARDIOMETABOLIC DISEASE

The essential component of DBCD is insulin resistance. When insulin secretion is unable to compensate for insulin resistance, the result is hyperglycemia and eventual prediabetes, identified on the basis of impaired FPG or impaired 2-hour postchallenge plasma glucose during an oral glucose tolerance test (40). Since up to 70% of patients with prediabetes have a lifetime risk of converting to T2D (40,41) (74% of individuals at age 45 [42]), primary prevention of T2D is paramount. Also, since those with prediabetes have significant CVD risk factors (36.6% with hypertension, 51.2% dyslipidemia, 24.3% tobacco use, and 5 to 7% 10-year cardiovascular event risk), secondary prevention is paramount by using effective diagnostic tests and treatments to reduce these risks and prevent CVD as early as possible (12,43).

There are three aspects to reducing the impact of prediabetes using primary and secondary prevention strategies. The first aspect is to prevent T2D. Progression to T2D can be prevented with interventions that improve insulin sensitivity, such as weight loss, healthy eating patterns, regular and sustained physical activity, and/or the use of diabetes medications (e.g., metformin, thiazolidinediones, and incretin-based therapies) (37,44-47). Left unchecked, prediabetes will progress to T2D in a majority of patients, while some others will continue with prediabetes or revert to normal glucose tolerance (48,49). While normoglycemia may be temporary with ultimate return of progressive glucose intolerance, individuals who revert to normal glucose tolerance are at diminished risk of T2D compared to those with stable prediabetes but remain at greater risk of T2D compared with those entirely without DBCD, particularly if there are other metabolic risk factors present or if they meet criteria for MetS (48,49). The second aspect is to prevent CVD. Prediabetes represents a state of clustered CVD risk factors, accelerated atherosclerosis, and increased risk for CVD events. Epidemiologically, elevated postprandial plasma glucose levels are more associated with increased CVD risk than FPG levels, particularly in women (50). Aggressive management of CVD risk factors is warranted in patients with prediabetes, though the exact degree and nature of intervention, as well as the subpopulations best served by this approach, remain unclear (38). The third aspect is to prevent T2D-related complications. The degree of dysglycemia in prediabetes is sufficient to cause microvascular complications of diabetes in some patients, as demonstrated in the Diabetes Prevention Program, where up to 10% of patients developed background retinopathy or neuropathy (51). Thus, in addition to CVD risk factor management, improvement of glycemic status in the prediabetes stage should be a specific target, though again, the exact degree and nature of improvement remains unclear.

Insulin resistance in association with overweight/obesity gives rise to MetS, which may or may not be accompanied by prediabetes (23). In this way, the natural history of DBCD intersects the natural history of ABCD at the level of insulin resistance, which involves abnormalities in the amount, distribution, and/or function of adipose tissue (35). The abnormal adipocyte secretome can affect the cardiometabolic nexus of disease manifestations, including prediabetes, MetS, nonalcoholic fatty liver disease, T2D, and CVD (52). In these patients, there is an infiltration of inflammatory macrophages in adipose tissue; dysregulated secretion of adipokines; impaired lipid storage leading to redistribution of lipid to the intra-abdominal compartment, liver, and muscle cells; and an exacerbation of insulin resistance (53-55). Hence, in those patients who develop prediabetes, ABCD is essentially indistinguishable from DBCD. However, not all patients with ABCD have DBCD. ABCD can also involve an augmentation in

adipose tissue mass without abnormalities in adipose tissue function. These patients are insulin sensitive with little or no increase in risk for T2D and CVD (55-58); however, the increase in adipose mass predisposes to biomechanical complications such as osteoarthritis, sleep apnea, urinary stress incontinence, and dysmobility/disability, in addition to some forms of cancer. Weight gain on an insulin-resistant background can exacerbate adipose tissue dysfunction and the cardiometabolic disease process (56-58), producing MetS and/or prediabetes and leading to T2D. However, weight gain on an insulin-sensitive background may confer minimal or no increase in risk for T2D and CVD, at least for the short term (56-58), but may still predispose to the biomechanical complications of ABCD on the basis of increased adipose tissue mass.

THE SPECIFIC CONTEXT OF DBCD AS THE INSULIN RESISTANCE-PREDIABETES-T2D SPECTRUM

Dysglycemia is loosely defined as any abnormality in glycemic status that is associated with disease, or the potential for disease, with the earliest citation in PubMed in 1951 (59). There is general consensus that both insulin resistance and pancreatic β -cell dysfunction are the key initiating physiologic events for development of T2D, conferred by complex interactions among genetic mutations and/or polymorphisms on a genomic scale, with epigenomic modifications and environmental factors (60). Recently, the role of pancreatic α -cell dysfunction and hyperglucagonemia has been investigated, wherein α -cells are stimulated by glucose-dependent insulinotropic polypeptide (GIP) and glucagon-like peptide 1 (GLP-1), and high glucagon levels stimulate hepatic glucose production (61). In fact, α -cells can influence pancreatic β -cell function by producing paracrine GLP-1 from proglucagon (62). Incretin resistance may also be a significant element of dysglycemia. This is evidenced by improvements in glycemic status after Roux-en-Y gastric bypass or biliopancreatic diversion, which are associated with sufficient rises in incretin levels that could overcome incretin resistance (63). Furthermore, reciprocal effects of β -cell function can reduce the number of α -cells, the relative α -cell/ β -cell distribution, and α -cell function (64).

At a molecular level, mitochondrial DNA methylation markers are associated with early stage (decreased insulin sensitivity) but not late-stage (abnormal FPG or A1C) dysglycemia (65). The pathophysiology of T2D is even more complex when the results of Lawlor et al (66) are considered, in which 248, 138, and 24 genes were identified that have differential expression in T2D versus nondiabetes from human β -, α -, and δ -cell single-cell transcriptomes, respectively.

The association of prediabetes with macrovascular disease, namely CVD, is well established among many

different patient populations and ethnicities (44,67-77). In the Maastricht Study, Sorenson et al (78) found that prediabetes is also independently associated with microvascular dysfunction, which may contribute to T2D-associated CVD. This microvascular disease association is due to hyperglycemia, which is often independent of the cardiovascular risk scores (78). Quantitatively, the effects on CVD risk of impaired fasting glucose, impaired glucose tolerance, and insulin resistance as part of the MetS vary in degree (23,79,80). Specifically, in a meta-analysis of prospective cohort studies, Huang et al (43) found that for those patients with prediabetes (defined by various impaired fasting glucose and impaired glucose tolerance criteria), the range of relative risks for composite CVD was 1.13 to 1.30, coronary heart disease 1.10 to 1.20, stroke 1.06 to 1.20, and all-cause mortality 1.13 to 1.32. Interestingly, in the Whitehall II Study, the incidence rate for major CVD events was greater for those patients with prediabetes diagnosed by A1C compared with either FPG or 2-hour postchallenge glucose (81).

Even so, the impact and relevance of prediabetes on CVD risk are supported by progression of diabetes-related complications without intervention (82-85), reduction of cardiometabolic risks with lifestyle and weight-loss interventions (44,86-92), and reduction in health care costs with interventions (93-99). In short, the question here is not whether glycemia-targeted interventions mitigate the

implied prediabetes-CVD causal association, but rather that recognition of the need to treat in earlier DBCD stages with comprehensive risk reduction strategies, namely, structured lifestyle interventions and weight-loss therapy, produces clinically significant outcome benefits (i.e., decreased incidence of T2D, MetS, and CVD).

The causality direction and underlying mechanism for the adiposity and prediabetes relationship continue to be elucidated. However, in support of the ABCD model, visceral (reflected by waist circumference and waist-to-hip ratio measurements) and intrahepatic adiposity are more strongly associated with prediabetes and T2D than general measures of adiposity (e.g., body mass index [BMI] and truncal fat) (100,101). Moreover, patients with prediabetes and overweight/obesity have more endothelial dysfunction (using the relative hyperemia index) than their normoglycemic, desirable-weight controls (102), consistent with a model of CVD risk involving complex interactions among adiposity, dysglycemia, and inflammation.

Interventions shown to be effective in preventing progression from prediabetes to T2D, such as exercise and weight loss, are well known to increase insulin sensitivity. The function of β -cells is altered in DBCD, with increases in the early stages of the disease to compensate for insulin resistance, followed by decreases that involve defects in glucose sensing, islet cell size, and insulin content in those individuals destined to progress to overt T2D (103-




Fig. 1. Dysglycemia-based chronic disease and the insulin resistance-prediabetes-type 2 diabetes (T2D) spectrum. Prediabetes is Stage 2 in the dysglycemia spectrum, or dysglycemia-based chronic disease (i.e., DBCD) framework. Key mechanistic drivers are strong independent predictors of a particular outcome in a complex system (117). Primordial prevention—to decrease diabetes risk factors in the general population; primary prevention—to decrease conversion to T2D in patients with prediabetes; secondary prevention—to detect disease and decrease morbidity in patients with early asymptomatic T2D; tertiary prevention—to decrease progression of T2D related complications; and quaternary prevention—to decrease overmanagement/-medicalization of any patient along the spectrum. Each dysglycemia stage can be viewed as the basis, or low or high risk, for chronic cardiovascular disease along a cardiometabolic spectrum.

110). β -Cell dysfunction is particularly pronounced in certain ethnic populations with higher prevalence rates of prediabetes and T2D, such as Asian Indians (111). Mechanisms related to these changes are being clarified, for instance by studying *db/db* mice with high FPG but normal postchallenge plasma glucose levels, in which the exocytic machinery is altered (112). Importantly, exercise and weight loss improve β -cell function in patients with prediabetes (113). Also, in patients with prediabetes and overweight/obesity, treatment with liraglutide (1.8 mg/day) or other GLP-1 receptor agonists is associated with an increased insulin secretion rate and β -cell sensitivity to glucose (114,115).

While DBCD exists along a continuum, it is useful to identify distinct stages as the disease progresses to formulate an actionable preventive care plan. This model is designed as a starting point for a more comprehensive approach to T2D and CVD by recognizing that these outcomes are the culmination of a common and protracted disease process. The natural history and clinical progression of DBCD indicate that there are opportunities for primordial (reducing risk factors in populations), primary, secondary, tertiary, and even quaternary (reducing overmedicalization) prevention. While AACE (30) and the American Diabetes Association (41) have issued position statements regarding prediabetes, no medications have as yet been approved for this indication, and it has been controversial as to whether prediabetes should merit therapy as a disease entity (36,116). The proposed evidence-based 4-stage model outlined below emphasizes the context of prediabetes as a component in the progression of DBCD, with opportunities for various prevention modalities to reduce evolution to T2D, CVD, or both (Fig. 1) (117).

1. **Stage 1 DBCD “Insulin Resistance”:** starts at birth with genomic/epigenomic risk determinants for insulin resistance. Molecular risk for islet-cell dysfunction under chronic stress of insulin resistance may also be present. Specific molecular screening or aggressive case finding tests to detect risk are not yet commercially available. However, a significant family history of T2D, *in utero* exposure to gestational diabetes, detection of hyperinsulinemia, or abdominal obesity (i.e., ABCD characterized by abnormalities in fat distribution) are easily obtained markers for this earliest DBCD stage and can therefore prompt action. Those with insulin resistance are at higher risk of progressing along the DBCD continuum and merit primary disease prevention: structured lifestyle interventions that reduce cardiometabolic risk factors and later DBCD stage likelihoods, respectively. A molecular roadmap for this process was recently presented by Jin et al (118), who identified a dynamic driver network

governing the key predisease-T2D transition mechanisms. It is postulated that appropriate interventions (such as structured lifestyle change) at these critical time points could prevent or delay progression of disease (118). Along these lines, Khara et al (119) found that in patients with a high genetic (molecular) risk for coronary heart disease, a favorable (healthy) lifestyle was associated with a nearly 50% reduction in relative risk. Pragmatically, all individuals can benefit from population-based primordial disease prevention given the pervasive diabetogenic and obesogenic environment. Primordial prevention strategies include health messaging and public education that create a culture of wellness, in addition to legislation on local, state, and federal levels that transition unhealthy to healthy built environments and target obstetrical care to ameliorate the *in utero* environment and mitigate intergenerational disease transmission.

2. **Stage 2 DBCD “Prediabetes”:** the emergence of detectable biochemical elevations in blood glucose with or without CVD risk factors or MetS traits. Screening and/or aggressive case finding for dysglycemia are critical in this stage. Prediabetes is associated with a high risk of future T2D due to the combination of insulin resistance and islet-cell dysfunction and for CVD as a result of accelerated atherosclerosis. Population-based primordial prevention strategies are applicable in this stage. Primary prevention strategies are indicated for these individuals, such as structured lifestyle modifications resulting in weight loss and enhanced physical fitness, that improve metabolic and CVD risk factors and prevent progression to Stage 3 DBCD, or T2D (47,120-127). At present, *prediabetes* is still considered as a predisease, but with further scientific delineation, prediabetes may eventually meet criteria for a true disease state. Moreover, secondary prevention strategies are indicated for prediabetes incorporating early detection of T2D, as well as interventions to reduce the progression of CVD. Pragmatically, successful interventions (e.g., lifestyle alone versus lifestyle + pharmacotherapy) at Stage 2 DBCD Prediabetes can depend on risks for CVD, T2D, and other T2D complications. Cardiometabolic Disease Staging (55) can effectively stratify risk for future T2D. Alternatively, patients may be segregated into subtypes based on various FPG and/or A1C cutoffs (e.g., 110 mg/dL [128] and 6.0% [42 mmol/mol] [129], respectively):
 - Stage 2a: low-risk prediabetes (e.g., FPG <110 mg/dL and/or A1C <6.0% [42 mmol/

- mol]) managed with intensive lifestyle intervention alone, versus
- Stage 2b: high-risk prediabetes (e.g., FPG ≥110 mg/dL and/or A1C ≥6.0%) managed with intensive lifestyle intervention and possibly pharmacotherapy (which may be

- no different than those modalities used to manage Stage 3 DBCD T2D and reduce risks for disease progression (e.g., blood pressure and lipid management [32]).
- 3. **Stage 3 DBCD “T2D”:** marked by biochemical elevations in blood glucose associated with statis-

Table 1
Dysglycemia-Based Chronic Disease Stages by Clinical Context, Pragmatic Relevance and Actions, and Evidence Base^a

DBCD stage	Clinical context	Pragmatic relevance and actions	Evidence base references
Stage I Molecular (Insulin Resistance)	(Epi)genetic Risk	Family history	131,132
	ABCD	Investigational molecular markers	133
	CMDS	Basal hyperinsulinemia	134
		Abnormal adiposity	23,35,55
		Policy/laws/education for a healthy built environment	135,136
		Lifestyle change	137,138
		Cardiometabolic health	139
Stage 2 Biochemical Cardiometabolic Risk (Prediabetes)	Abnormal IFG, IGT, A1C	Screen/manage other MetS components	23
	GDM	ILI	31,44,47,90,92,121-127,140
	PCOS	Consider pharmacological therapy with Stage 2b	31,141
		Cardiometabolic risk	23
Stage 3 Biochemical Disease (Type 2 Diabetes)	Abnormal IFG, IGT, A1C	ILI	26
		Pharmacological therapy if needed for glycemic target	26
		Cardiometabolic disease	23
Stage 4 Vascular Complications (Type 2 Diabetes with Complications)	T2D complications	ILI	26
		Pharmacological therapy for glycemic targets	26
		Pharmacological/procedural therapy for specific complications	26
		Consider procedure with risk factors to achieve glycemic targets and complication management	142
		Cardiometabolic event biomarkers	143

Abbreviations: A1C = hemoglobin A1c; ABCD = adiposity-based chronic disease; CMDS = cardiometabolic disease staging system; DBCD = dysglycemia-based chronic disease; IFG = impaired fasting glucose; GDM = gestational diabetes; IGT = impaired glucose tolerance; ILI = intensive lifestyle intervention; MetS = metabolic syndrome; PCOS = polycystic ovary syndrome; T2D = type 2 diabetes.

^aThe severity of T2D complications is provided using the CMDS and other published information. Intensive lifestyle intervention consists of weight loss and strength training (when needed to optimize body composition), physical activity, healthy eating patterns, improved sleep hygiene, stress reduction, tobacco cessation, etc. Pharmacologic agents associated with T2D risk reduction are: phentermine/topiramate (49-89%); metformin (26-77%); troglitazone (54-75%); pioglitazone (70%); rosiglitazone (55%); orlistat (45%); lorcaserin (38%); and acarbose (25%) (141).

tically increased risk for T2D-related complications amenable to secondary prevention strategies. Aggressive case finding for CVD and T2D complications is critical in this stage. In patients with T2D with nonvascular complications, tertiary prevention strategies are needed to prevent further morbidity. Population-based primordial prevention strategies, used to reduce risk in the general population, are also applicable in this stage.

4. **Stage 4 DBCD “Vascular Complications”:** defined by the clinical presence of T2D microvascular complications (e.g., retinopathy, nephropathy, and neuropathy) and/or T2D macrovascular disease events (e.g., myocardial infarction, stroke, amputation, and ischemic foot ulcer), representing end-stage DBCD. These forms of organ dysfunction require tertiary prevention, or treatment, to ameliorate or reduce further deterioration. In this process, complications are diagnosed and treatment approaches fashioned according to mortality risks:
- Stage 4a: mild complications,
 - Stage 4b: moderate complications, and
 - Stage 4c: severe complications (130).

Population-based primordial prevention strategies are also applicable for stage 4 DBCD and can help reduce the emergence of new CVD risk factors.

Rather than using a continuous spectrum with blurred demarcations among pathologic states, the above discrete stages are parsed out for practical application with respect to clinical context, pragmatic relevance, and an evidence base (Table 1) (131-143). Each of the stages is also amenable to quaternary prevention, in which overmedicalization and iatrogenesis are minimized.

DYSGLYCEMIA-BASED CHRONIC DISEASE— A NEW MULTIMORBIDITY T2D MODEL

Chronic disease is generally regarded as a pathological condition lasting for more than 3 months (144). In order for a medical condition to be viewed within a preventive care model, 3 criteria should exist: (1) association with a significant amount of human suffering; (2) safe and effective, affordable, and validated screening tools; and (3) safe and effective, affordable, and validated interventions for each prevention type (145,146). Even when there is a single etiology, there are many downstream paths that branch out producing multiple sequelae, complications, and morbid events. With complex interactions occurring over varying time scales, a steady state can emerge that characterizes the multimorbid chronic disease state (144). This model becomes even more complicated when there are multiple initiating, interacting, and/or concurrent etiologies, such as

T2D, obesity, coronary artery disease, cancer, depression, inflammation, frailty, and/or medications. The complexity of chronic disease emphasizes the importance of psychosocial, economic, and behavioral interventions, research metrics related to patient-defined goals of care, community-based leadership, and service- and therapeutic-level integration (147-149). The model will also need to be context-adaptable for people of different cultures/ethnicities (e.g., prediabetes to T2D conversion rates are faster for patients of South Asian > African > European ancestries; and in Asians, β -cell dysfunction predominates over insulin resistance [149-153]), as well as socio-economic strata and ability/disability (e.g., using patient-centered medical homes to improve diabetes self-management education and support [154]). In a study by Eisenstat et al (155) primarily addressing insulin use, patients with prediabetes and T2D were followed by a coordinated team in a network of primary care physicians, using evidence-based recommendations addressing medical, behavioral, and nutrition/lifestyle components. Certain factors were identified that facilitated care: diverse stakeholder engagement and buy-in with the theoretical basis of the model, institutional alignment of priorities, and credits for participation and implementation (155). The close integration of pharmacists can also improve adherence factors in the care model (156).

Overall, a chronic care model incorporates the following elements:

- a prepared and informed patient,
- capable, knowledgeable, and skilled health care professionals,
- organization of health care services,
- self-care support,
- clinical decision support,
- clinical information systems,
- design of the service delivery system, and
- community resources (157,158).

According to a systematic review of randomized, controlled trials involving patients with T2D from the Medline and Cochrane Library electronic databases, Baptista et al (157) found that the use of isolated chronic care model elements may be insufficient to improve clinical outcomes, but incorporating all of the elements together may prove beneficial. Figure 2 depicts the AACE DBCD multimorbidity T2D chronic care model based on the template used for the AACE obesity chronic care model that incorporates these and other elements (33). Each component is important in this preventive paradigm, which may have the greatest potential in terms of prevention of morbidity and mortality when patients are actively engaged in intervention at stage 2 DBCD Prediabetes.

The AACE DBCD model will require both clinical and economic validation. Research will need to address each component with respect to establishing thresholds




Fig. 2. American Association of Clinical Endocrinologists/American College of Endocrinology dysglycemia-based chronic disease multimorbidity type 2 diabetes care model. The built environment contextualizes diabetes care and consists of human-made entities. Decision support systems include electronic implementations of clinical practice guidelines. Delivery systems designs involve multidisciplinary care teams for glycemic control as well as complications. Empanelment is the process of linking patients with a diabetes care team and primary care physician, as well as serving as basis for performance metrics.

(molecular, biochemical, and clinical) for engagement and action, adhering with evidence-based management protocols appropriate for primary, secondary, and tertiary prevention, and incorporating suitable clinical outcome metrics. For instance, can transitions among the four stages and additional substages in the DCBD model be better demarcated using novel CGM technologies, cardiometabolic biomarkers, and metabolite/metabolomic biomarkers, along with revised fasting and postchallenge plasma glucose levels, judicious A1C testing, and self-monitoring of plasma glucose (132,159-161)? Can a FPG cutoff at 110 mg/dL (instead of 100 mg/dL) and/or A1C cutoff at 6.0% (42 mmol/mol) (instead of 5.7% [39 mmol/mol]) be used to discriminate a low-risk stage 2a (managed with lifestyle change) from a high-risk stage 2b (managed with lifestyle change and, if needed, pharmaceuticals) be validated (27,128,129,162)? The points here are that more research and analysis are needed to optimize these classifiers (163), rationally reduce unnecessary testing and pharmacotherapy (164), and improve lifestyle interventions for all subsets of the general population. Given the high prevalence of DBCD, risk stratification will be critically important from both the perspective of risk-benefit and cost-effectiveness in bringing more aggressive therapies to those patients at highest risk at each stage of preven-

tion/treatment. For example, using the Cardiometabolic Disease Staging system—a T2D risk stratification strategy employing the number of MetS traits and presence of prediabetes—demonstrates that a medicine-assisted weight-loss intervention can prevent T2D in high-risk individuals with lower numbers-needed-to-treat than in low-risk individuals (55,165,166).

Various economic modeling methods have already demonstrated positive impact for chronic disease management (167,168), including T2D (169). Introducing communications technology has been shown to facilitate adoption of a diabetes chronic care model and use of self-management strategies, while also lowering costs (170). Future economic studies on T2D prevention will need to concentrate on context, scope, payment models, income levels, self-management support, and impact on the labor market (169-172). The economic model will also need validation for patients of different cultures and ethnicities, both of which are associated with differences in key mechanistic drivers, phenotypic expression of disease, and conversion rates among the DCBD stages (173-176).

Currently, treatment of DBCD often begins at the time of T2D diagnosis. The DBCD paradigm places insulin resistance, prediabetes, MetS, T2D, and CVD within the context of a single chronic disease based on scientific

data and a current understanding of common pathophysiologic mechanisms. The clinical implications are that initiation of therapy only at the time of Stage 3 DBCD T2D or Stage 4 DBCD Vascular Complications diagnosis is not an optimal care approach. Rather, initiating DBCD management at earlier Stage 1 DBCD Insulin Resistance or Stage 2 DBCD Prediabetes can more effectively reduce disease progression, symptom burden, unhealthy population metrics, and overall health care costs. In fact, there is considerable scientific evidence that validate earlier intervention in high-risk individuals, including prediabetes, to reduce CVD (177-180). This new paradigm brings into focus health policy, the built environment, lifestyle medicine, and a philosophical position to restrain overmedicalization. With the perspective that T2D is only one stage in a progressive chronic disease process, a complement of preventive care modalities could rationally be initiated at any stage in DBCD progression depending upon risk-benefit and overall impact on public health, disease burden, and quality of life.

CONCLUSION

Recognizing and managing prediabetes is a necessary component for an effective personalized and population-based T2D care plan. In order to substantiate this position, AACE has formulated a DBCD multimorbidity care model consisting of four distinct stages in the general context of ABCD and cardiometabolic health and specific context along the insulin resistance-prediabetes-T2D spectrum that are actionable in a preventive care paradigm. Thus, prediabetes is regarded as stage 2 in the DBCD framework. This re-interpretation of prediabetes is consistent with other approaches, such as the recent move away from a strict glucocentric definition of prediabetes (181) to the AACE complications-centric (instead of BMI-centric) framework for obesity care (33,35).

The controversy over whether there is value, cost-effectiveness, or clinical benefit in diagnosing and/or managing the prediabetes state is resolved by regarding the problem, not in isolation where the evidentiary basis of

prediabetes recognition and management are challenged, but rather as part of comprehensive attention to a progressive, prevalent, and impactful metabolic disease, on a host of scales, managed by multiple concurrent preventive medicine strategies. This AACE position is consistent with a portfolio of endocrine disease care models that prioritize patient-centered care, evidence-based medicine, care models for complex and multimorbid chronic disease, the current health care environment, and a societal mandate for high value and good health.

With the refinement of the ABCD and DBCD diagnostic terms and care models, AACE will continue to develop comparable activities to eventually construct an actionable global cardiometabolic-based chronic disease framework. For instance, just as ABCD and DBCD incorporate and are part of inflammatory networks, new models of blood pressure homeostasis incorporate polarization of interstitial macrophage populations, direct effects on sodium sequestration, and networking effects on renovascular mechanisms of hypertension (52,182). Hence, chronic disease models of adiposity, dysglycemia, dyslipidemia, hypertension, and inflammation converge onto a unified theory of cardiometabolic risk. Consequently, any controversy over whether prediabetes exists or not will be obviated within this larger framework purposed to prevent morbidity and mortality from composite cardiometabolic risks. Moreover, at some point in the future, the concept of lifetime T2D risk can be formulated based on DBCD prevalence rates to inform decision-making and gauge whether diabetes fate can be avoided or merely delayed. At present, the DBCD framework is not applicable to type 1 diabetes.

Tactically, as with the AACE ABCD framework, a new diagnostic coding system can be engineered for prediabetes and T2D within the DBCD framework that can translate these concepts into organized action. As with ABCD, one can envision a robust coding system cross-tabulating each DBCD stage with specific phenotypes and complications. With validation and the development of appropriate reimbursement structures, a distinct transformation in endocrine health care can be realized.

Glossary Term

Adiposity

Definition (glossary terms in bold)

related to the amount, distribution, or function of fat tissue, cells containing fat, and/or adipocytes

Adiposity-based chronic disease

a chronic disease state resulting from abnormal **adiposity**

Aggressive case finding

process of identifying persons most likely to have a **chronic disease** based on presence of risk factors, manifestations, and likely benefit from treatment

Alpha (α)-cell dysfunction

mechanisms related to abnormal glucagon secretion in pancreatic α -cells in the context of increased hepatic glucose production, regulation by GIP and GLP-1, and **type 2 diabetes** pathogenesis and natural history

Beta (β)-cell dysfunction

mechanisms related to abnormal insulin secretion in pancreatic β -cells in the context of **type 2 diabetes** pathogenesis and natural history

Biochemical risk	abnormal blood tests that are correlated with an increased risk for disease (e.g., specified fasting plasma glucose, 2-hour post challenge plasma glucose, or hemoglobin A1c levels that diagnose prediabetes)
Built environment	the human-made world, including buildings and other structures, sidewalks and running paths, elevators and stairs, and all other aspects of human-made surroundings for daily life; this does not include natural surroundings (e.g., climate, geography, or terrain) or culture (e.g., religion, socio-economics, or politics)
Cardiometabolic risk	term that encompasses both metabolic risk for diabetes and vascular risk for coronary heart disease and/or stroke due to common pathophysiologic processes. This can also be understood as a broader and more interpretative term that describes the aggregate risk for atherosclerotic cardiovascular disease based on component metabolic factors, such as adiposity and dysglycemia
Cardiovascular disease	diseases of the heart and/or blood vessels that include: coronary heart, rheumatic heart, congenital heart, cerebrovascular, and peripheral arterial disease, as well as deep vein thrombosis, pulmonary embolus, heart failure, arrhythmia, and other valvular problems
Chronic disease	a disease state that persists for 3 months or longer; generally associated with a complex array of predisposing biological, environmental, and cultural factors, clinical manifestations, and complication risks
Diabetes-related complications	end-organ dysfunction that results from diabetes (e.g., nephropathy, retinopathy, and neuropathy)
Diabetes (mellitus)	a chronic disease state characterized by abnormally high blood glucose levels and increased risk for premature mortality and diabetes-related complications , such as retinopathy and cardiovascular disease
Disease	a diagnosed medical condition that impairs normal function, has characteristic signs or symptoms, and causes harm or morbidity. At present, diabetes is considered a disease , but prediabetes is considered a predisease . Diseases are managed with secondary prevention or tertiary prevention strategies.
Dysglycemia	a more general term that includes all forms of diabetes (usually type 2 diabetes) and prediabetes , including states of increased molecular risk for diabetes
Dysglycemia-based chronic disease	a chronic disease state resulting from dysglycemia
Hyperglycemia	elevated plasma glucose, generally interpreted in a specific physiologic context, such as fasting, postprandial, postchallenge , stress, or pregnancy
Incretin resistance	a pathophysiologic state in which β -cells exhibit subnormal or absent insulin secretory responses to incretins and α -cells exhibit partial or unsuppressed glucagon secretory responses to incretins
Insulin resistance	a pathophysiologic state in which cells exhibit subnormal responses to insulin; generally understood in the context of hepatocytes and myocytes in type 2 diabetes , as an etiologic event producing hyperglycemia , exacerbated by hyperglucagonemia and α-cell dysfunction , and either independent or dependent on β-cell dysfunction
Intervention	an action that encompasses all types of prevention
Lifestyle	one's manner of living, generally consisting of attitudes, beliefs, practices, culture, and interactions with the natural and built environment
Lifestyle medicine	the nonpharmacologic and nonsurgical/nonprocedural management of chronic disease
Metabolic syndrome	cluster of specific and residual risks for cardiovascular disease ; depending on the consensus definition, specific risks include insulin resistance , obesity , hypertension, hypertriglyceridemia, and markers of vascular inflammation and thrombosis
Molecular Risk	a predisease state characterized by the association of statistical- or network-based genetic or molecular markers (classifiers) with an increased incidence or prevalence rate of a chronic disease state
Obesity	a term used to describe the chronic disease state associated with a body mass index at or over specific ethnicity-adjusted cutoffs (e.g., 30 kg/m ² for Caucasians) presumed to be due to an increased amount of adiposity
Overweight	a term used to describe the pre- obesity state associated with a body mass index at or over specific ethnicity-adjusted cutoffs (e.g., 25 kg/m ² for Caucasians) presumed to be due to an increased amount of adiposity

Postchallenge	describes the context for hyperglycemia when a plasma glucose level is measured a specified time after administration of a specified amount of glucose (e.g., 75 g of oral glucose and measuring plasma glucose 2 hours later; the “2-hour oral glucose tolerance test”)
Prediabetes	a predisease state characterized by one or more abnormal biochemical tests (e.g., fasting plasma glucose, postchallenge plasma glucose, or hemoglobin A1c) that are associated with increased incidence or prevalence of type 2 diabetes and cardiovascular disease
Predisease	a term describing an early stage of chronic disease , before the disease is diagnosed, and during which time screening or aggressive case finding may be performed, and primary prevention strategies implemented
Primary prevention	population- and individual-based strategies applied before a disease is diagnosed in order to prevent disease occurrence; this includes effective screening and aggressive case finding , primordial prevention tactics, and in some cases, the judicious use of pharmacotherapy
Primordial prevention	population-based strategies applied at all stages along the chronic disease spectrum to reduce the incidence and prevalence of risk factors for chronic disease(s) ; examples include beneficial changes in behavior, the built environment , socio-economics, and health care policy
Quaternary prevention	population- and individual-based strategies applied to all stages along the chronic disease spectrum to reduce over- medicalization; specifically decreasing adverse events, improving value, and encouraging ethical decision-making; emphasizes primordial , primary , and secondary prevention education and practice
Residual risk	the amount of risk for a chronic disease that remains once specific, known component risks have been reduced (e.g., the risk for cardiovascular disease after hypertension, obesity , insulin resistance , and dyslipidemia risks have been controlled in patients with metabolic syndrome)
Screening	process of identifying persons most likely to have a common chronic disease but without clinical risks or manifestations, based on the application of a test, where safe and cost-effective treatments are available
Secondary prevention	population- and individual-based strategy applied to diagnosed chronic disease states in early stages to reduce the general impact or morbidity of the disease ; modalities generally include regular medical examinations, effective diagnostic testing, lifestyle medicine , and when needed, pharmacotherapy
Tertiary prevention	population- and individual-based strategy applied to diagnosed chronic disease states in later stages to reduce the impact of specific symptoms and slow the progression of specific complications; this can be the costliest and most prevalent prevention modality, including relatively expensive pharmacotherapy and procedures, and is the principal target of quaternary prevention
Type 2 diabetes	a specific type of diabetes that results from α-cell dysfunction , β-cell dysfunction , and insulin resistance , generally characterized by insufficient insulin action to prevent hyperglycemia but sufficient insulin action to prevent ketoacidosis, except in later stages and/or atypical forms; associated with specific diabetes-related complications

DISCLOSURES

Dr. Mechanick received honoraria for lectures and program development from Abbott Nutrition. Dr. Garber is a consultant to Novo Nordisk and Intarcia Therapeutics. Dr. Grunberger reports research contracts with Novo Nordisk and Medtronic and is on the speakers bureau of Eli Lilly, Novo Nordisk, Sanofi, Janssen, Boehringer Ingelheim, and AstraZeneca. Dr. Handlesman reports research grants and consultant and speaker honoraria from Aegerion, Amgen, AstraZeneca, Bristol-Myers-Squibb, Boehringer Ingelheim, BI-Lilly, Gan & Lee, Gilead, Grifols, Hamni, Intarcia, Janssen, Lexicon, Lilly, Merck, Mylan, Merck-Pfizer, Novo Nordisk, Regeneron, and Sanofi. Dr. Garvey

reports that he is on the advisory boards of Novo Nordisk, Janssen, Astra Zeneca, Alexion, Merck, American Medical Group Association, National Diabetes and Obesity Research Institute; he has received research funding from Merck, Astra Zeneca, Weight Watchers, Eisai, Sanofi, Pfizer, Novo Nordisk, Lexicon, Elcelyx; and he has stock ownership (publically traded) in Eli Lilly, Pfizer, Novartis, Merck, Isis, Bristol-Myers-Squibb, and Affymetrix.

REFERENCES

1. **Hegazi R, El-Gamal M, Abdel-Hady N, Hamdy O.** Epidemiology of an risk factors for type 2 diabetes in Egypt. *Ann Global Health.* 2015;81:814-820.

2. **John H.** Glucose tolerance and its value in diagnosis. *J Metabolic Res.* 1922;1:497-548.
3. **Mosenthal HO, Barry E.** Criteria for and interpretation of normal glucose tolerance tests. *Ann Intern Med.* 1950;33:1175-1194.
4. **McCartney P, Keen H, Jarrett RJ.** The Bedford Survey: observations on retina and lens of subjects with impaired glucose tolerance and in controls with normal glucose tolerance. *Diabetes Metab.* 1983;9:303-305.
5. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care.* 1997;20:1183-1197.
6. AACE Diabetes Scientific Advisory Group. American Association of Clinical Endocrinologists/American College of Endocrinology statement on the use of hemoglobin A1c for the diagnosis of diabetes. *Endocr Pract.* 2010;16:155-156.
7. **Dorcely B, Katz K, Jagannathan R, et al.** Novel biomarkers for prediabetes, diabetes, and associated complications. *Diab Met Syndr Obes.* 2017;10:345-361.
8. **Fonseca VA, Grunberger G, Anhalt H, et al.** Continuous glucose monitoring: a consensus conference of the American Association of Clinical Endocrinologists and American College of Endocrinology. *Endocr Pract.* 2016;22:1008-1021.
9. **Kaul N, Ali S.** Genes, genetics, and environment in type 2 diabetes: implication in personalized medicine. *DNA Cell Biol.* 2016; 35:1-12.
10. Centers for Disease Control and Prevention. 2014 National Diabetes Statistics Report. Available at: <https://www.cdc.gov/diabetes/pdfs/data/2014-report-estimates-of-diabetes-and-its-burden-in-the-united-states.pdf>. Accessed September 17, 2018.
11. Centers for Disease Control and Prevention. 2017 National Diabetes Statistics Report. Available at: <https://www.cdc.gov/diabetes/pdfs/data/statistics/national-diabetes-statistics-report.pdf>. Accessed October 28, 2017.
12. **Ali MK, Bullard KM, Saydah S, Imperatore G, Gregg EW.** Cardiovascular and renal burdens of prediabetes in the USA: analysis of data from serial cross-sectional surveys, 1988-2014. *Lancet Diabetes Endocrinol.* 2018;6:392-403.
13. **Dall TM, Yang W, Halder P, et al.** The economic burden of elevated blood glucose levels in 2012: diagnosed and undiagnosed diabetes, gestational diabetes mellitus, and prediabetes. *Diabetes Care.* 2014;37:3172-3179.
14. **Li G, Zhang P, Wang J, et al.** Cardiovascular mortality, all-cause mortality, and diabetes incidence after lifestyle intervention for people with impaired glucose tolerance in the Da Qing Diabetes Prevention Study: a 23-year follow-up study. *Lancet Diabetes Endocrinol.* 2014;2:474-480.
15. **Mudaliar U, Zabetian A, Goodman M, et al.** Cardiometabolic risk factor changes observed in diabetes prevention programs in US settings: a systematic review and meta-analysis. *PLoS Med.* 2016;13:e1002095.
16. **Uusitupa M, Peltonen M, Lindström J, et al.** Ten-year mortality and cardiovascular morbidity in the Finnish Diabetes Prevention Study—secondary analysis of the randomized trial. *PLoS One.* 2009;4:e5656.
17. **Balk EM, Earley A, Raman G, Avendano EA, Pittas AG, Remington PL.** Combined diet and physical activity promotion programs to prevent type 2 diabetes among persons at increased risk: a systematic review for the Community Preventive Services Task Force. *Ann Intern Med.* 2015;163:437-451.
18. **Jackson WP, Woolf N.** The natural history of prediabetes. *N Engl J Med.* 1956;255:1183-1185.
19. **Di Pino A, Urbano F, Piro S, Purello F, Rabuazzo AM.** Update on pre-diabetes: focus on diagnostic criteria and cardiovascular risk. *World J Diabetes.* 2016;7:423-432.
20. **Hanefeld M, Pistrosch F, Koehler C, Chiasson JL.** Conversion of IGT to type 2 diabetes mellitus is associated with incident cases of hypertension: a post-hoc analysis of the STOP-NIDDM trial. *J Hypertens.* 2012;30:1440-1443.
21. **Jagannathan R, Sevick MA, Fink D, et al.** The 1-hour post-load glucose level is more effective than HbA1c for screening dysglycemia. *Acta Diabetol.* 2016;53:543-550.
22. **Viera AJ.** Predisease: when does it make sense? *Epidemiol Rev.* 2011;33:122-134.
23. **Sperling LS, Mechanick JI, Neeland IJ, et al.** The CardioMetabolic Health Alliance: working toward a new care model for the metabolic syndrome. *J Am Coll Cardiol.* 2015;66:1050-1067.
24. **Diabetes Prevention Program Research Group.** Long-term effects of lifestyle intervention or metformin on diabetes development and microvascular complications over 15-year follow-up: the Diabetes Prevention Program Outcomes Study. *Lancet Diabetes Endocrinol.* 2015;3:866-875.
25. **Yudkin JS, Montori VM.** The epidemic of pre-diabetes: the medicine and the politics. *BMJ.* 2014;349:g4485.
26. **Reach G.** Simplistic and complex thought in medicine: the rationale for a person-centered care model as a medical revolution. *Patient Prefer Adherence.* 2016;10:449-457.
27. **Davidson M, Kahn RA.** A reappraisal of prediabetes. *J Clin Endocrinol Metab.* 2016;101:2628-2635.
28. **Einhorn D, Reaven GM, Cobin RH, et al.** ACE position statement on the insulin resistance syndrome. *Endocr Pract.* 2003;9:237-252.
29. **Mechanick JI, Cobin RH, Einhorn D, Handelsman Y, Hellman R, Jellinger PS.** American College of Endocrinology/American Association of Clinical Endocrinologists: reaffirmation of the 2003 ACE insulin resistance syndrome (IRS) position statement. Available at: <https://www.aace.com/files/position-statements/irsstatement.pdf>. Accessed January 2, 2017.
30. **Garber AJ, Handelsman Y, Einhorn D, et al.** Diagnosis and management of prediabetes in the continuum of hyperglycemia: when do the risks of diabetes begin? A consensus statement from the American College of Endocrinology and the American Association of Clinical Endocrinologists. *Endocr Pract.* 2008;14:933-946.
31. **Handelsman Y, Bloomgarden ZT, Grunberger G, et al.** American Association of Clinical Endocrinologists and American College of Endocrinology – clinical practice guidelines for developing a diabetes mellitus comprehensive care plan – 2015. *Endocr Pract.* 2015;21(suppl 1):1-87.
32. **Garber AJ, Abrahamson MJ, Barzilay JI, et al.** Consensus statement by the American Association of Clinical Endocrinologists and American College of Endocrinology on the comprehensive type 2 diabetes management algorithm – 2018 executive summary. *Endocr Pract.* 2018;24:91-120.
33. **Garvey WT, Mechanick JI, Brett EM, et al.** American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract.* 2016;22(suppl 3):1-203.
34. **Jellinger PS, Smith DA, Mehta AE, et al.** American Association of Clinical Endocrinologists' guidelines for management of dyslipidemia and prevention of atherosclerosis. *Endocr Pract.* 2012;18(suppl 1):1-78.
35. **Mechanick JI, Hurley DL, Garvey WT.** Adiposity-based chronic disease as a new diagnostic term: the American Association of Clinical Endocrinologists and American College of Endocrinology position statement. *Endocr Pract.* 2017;23:372-378.
36. **Cefalu WT, Buse JB, Tuomilehto J, et al.** Update and next steps for real-world translation of interventions for type 2 diabetes prevention: reflections from a diabetes care editors' expert forum. *Diabetes Care.* 2016;39:1186-1201.
37. **Grams J, Garvey WT.** Weight loss and the prevention and treatment of type 2 diabetes using lifestyle therapy, pharmacotherapy, and bariatric surgery: mechanisms of action. *Curr Obes Rep.* 2015;4:287-302.
38. **Ahlqvist E, Storm P, Käräjämäki A, et al.** Novel subgroups of adult-onset diabetes and their association with outcomes: a data-driven cluster analysis of six variables. *Lancet Diabetes Endocrinol.* 2018;6:361-369.
39. **Stefan N, Fritsche A, Schick F, Häring HU.** Phenotypes of prediabetes and stratification of cardiometabolic risk. *Lancet Diabetes Endocrinol.* 2016;4:789-798.
40. **Tabák AG, Herder C, Rathmann W, Brunner EJ, Kivimäki M.** Prediabetes: a high-risk state for diabetes development. *Lancet.* 2012;379:2279-2290.
41. **Nathan DM, Davidson MB, DeFronzo RA, et al.** American Diabetes Association. Impaired fasting glucose and impaired

- glucose tolerance: implications for care. *Diabetes Care.* 2007;30:753-759.
42. **Ligthart S, van Herpt TT, Leening MJ, et al.** Lifetime risk of developing impaired glucose metabolism and eventual progression from prediabetes to type 2 diabetes: a prospective cohort study. *Lancet Diabetes Endocrinol.* 2016;4:44-51.
 43. **Huang Y, Cai X, Mai W, Li M, Hu Y.** Association between prediabetes and risk of cardiovascular disease and all-cause mortality: systematic review and meta-analysis. *BMJ.* 2016;355:i5953.
 44. **Knowler WC, Barrett-Connor E, Fowler SE, et al.** Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med.* 2002;346:393-403.
 45. **Garvey WT, Ryan DH, Henry R, et al.** Prevention of type 2 diabetes in subjects with prediabetes and metabolic syndrome treated with phentermine and topiramate extended-release. *Diabetes Care.* 2014;37:912-921.
 46. **le Roux CW, Astrup A, Fujioka K, et al.** 3 years of liraglutide versus placebo for type 2 diabetes risk reduction and weight management in individuals with prediabetes: a randomised, double-blind trial. *Lancet.* 2017;389:1399-1409.
 47. **DREAM (Diabetes REDuction Assessment with ramipril and rosiglitazone Medication) Trial Investigators: Gerstein HC, Yusuf S, Bosch J, et al.** Effect of rosiglitazone on the frequency of diabetes in patients with impaired glucose tolerance or impaired fasting glucose: a randomised controlled trial. *Lancet.* 2006;368:1096-1105.
 48. **Perreault L, Pan Q, Mather KJ, Watson KE, Hamman RF, Kahn SE.** Effect of regression from prediabetes to normal glucose regulation on long-term reduction in diabetes risk: results from the Diabetes Prevention Program Outcomes Study. *Lancet.* 2012;379:2243-2251.
 49. **Herman WH, Pan Q, Edelstein SL, et al.** Impact of lifestyle and metformin interventions on the risk of progression to diabetes and regression to normal glucose regulation in overweight or obese people with impaired glucose regulation. *Diabetes Care.* 2017;40:1668-1677.
 50. **Cavalot F, Petrelli A, Traversa M, et al.** Postprandial blood glucose is a stronger predictor of cardiovascular events than fasting blood glucose in type 2 diabetes mellitus, particularly in women: lessons from the San Luigi Gonzaga Diabetes Study. *J Clin Endocrinol Metab.* 2006;91:813-819.
 51. **Diabetes Prevention Program Research Group.** The prevalence of retinopathy in impaired glucose tolerance and recent-onset diabetes in the Diabetes Prevention Program. *Diabet Med.* 2007;24:137-144.
 52. **Mechanick JI, Zhao S, Garvey WT.** The adipokine-cardiovascular-lifestyle network. *J Am Coll Cardiol.* 2016;68:1785-1803.
 53. **Olefsky JM, Glass CK.** Macrophages, inflammation, and insulin resistance. *Annu Rev Physiol.* 2010;72:219-246.
 54. **Lara-Castro C, Fu Y, Chung BH, Garvey WT.** Adiponectin and the metabolic syndrome: mechanisms mediating risk for metabolic and cardiovascular disease. *Curr Opin Lipidol.* 2007;18:263-270.
 55. **Guo F, Moellering DR, Garvey WT.** The progression of cardiometabolic disease: Validation of a new cardiometabolic disease staging system applicable to obesity. *Obesity (Silver Spring).* 2014;22:110-118.
 56. **Stefan N, Kantartzis K, Machann J, et al.** Identification and characterization of metabolically benign obesity in humans. *Arch Intern Med.* 2008;168:1609-1616.
 57. **Wildman RP, Muntner P, Reynolds K, et al.** The obese without cardiometabolic risk factor clustering and the normal weight with cardiometabolic risk factor clustering: prevalence and correlates of 2 phenotypes among the US population (NHANES 1999-2004). *Arch Intern Med.* 2008;168:1617-1624.
 58. **Guo F, Garvey WT.** Cardiometabolic disease risk in metabolically healthy and unhealthy obese: stability of metabolic health status in adults. *Obesity (Silver Spring).* 2016;24:516-525.
 59. **Soler B.** Double hyperglycemic test in the diagnosis of disorders of hydrocarbon metabolism (dysglycemia) [in undetermined language]. *Dia Med.* 1951;23:2997-3006.
 60. **Edwards CM, Cusi K.** Prediabetes: a worldwide epidemic. *Endocr Metab Clin North Am.* 2016;45:751-764.
 61. **Lund A, Bagger JI, Christensen M, Knop FK, Vilsbøll T.** Glucagon and type 2 diabetes: the return of the alpha cell. *Curr Diab Rep.* 2014;14:555.
 62. **Traub S, Meier DT, Schulze F, et al.** Pancreatic α cell-derived glucagon-related peptides are required for β cell adaptation and glucose homeostasis. *Cell Rep.* 2017;18:3192-3203.
 63. **Salinari S, Mingrone G, Bertuzzi A, Previtto E, Capristo E, Rubino F.** Downregulation of insulin sensitivity after oral glucose administration: evidence for the anti-incretin effect. *Diabetes.* 2017;66:2756-2763.
 64. **Henquin JC, Rahier J.** Pancreatic α cell mass in European subjects with type 2 diabetes. *Diabetologia.* 2011;54:1720-1725.
 65. **Zheng LD, Linarelli LE, Brooke J, et al.** Mitochondrial epigenetic changes link to increased diabetes risk and early-stage prediabetes indicator. *Oxid Med Cell Longev.* 2016;2016:5290638.
 66. **Lawlor N, George J, Bolisetty M, et al.** Single-cell transcriptomes identify human islet cell signatures and reveal cell-type-specific expression changes in type 2 diabetes. *Genome Res.* 2017;27:208-222.
 67. **Chiasson JL, Josse RG, Gomis R, Hanefeld M, Karasik A, Laasko M.** Acarbose treatment and the risk of cardiovascular disease and hypertension in patients with impaired glucose tolerance: the STOP-NIDDM trial. *JAMA.* 2003;290:486-494.
 68. **Levitzyk YS, Pencina MJ, D'Agostino RB, et al.** Impact of impaired fasting glucose on cardiovascular disease: the Framingham Heart Study. *J Am Coll Cardiol.* 2008;51:264-270.
 69. **Glucose tolerance and mortality: comparison of WHO and American Diabetes Association diagnostic criteria. The DECODE study group. European Diabetes Epidemiology Group. Diabetes epidemiology: collaborative analysis of diagnostic criteria in Europe. Lancet.** 1999;354:617-621.
 70. **Barr EL, Zimmet PZ, Welborn TA, et al.** Risk of cardiovascular and all-cause mortality in individuals with diabetes mellitus, impaired fasting glucose, and impaired glucose tolerance: the Australian Diabetes, Obesity, and Lifestyle Study (AusDiab). *Circulation.* 2007;116:151-157.
 71. **Muhlestein JB, Anderson JL, Horne BD, et al.** Intermountain Heart Collaborative Study Group. Effect of fasting glucose levels on mortality rate in patients with and without diabetes mellitus and coronary artery disease undergoing percutaneous coronary intervention. *Am Heart J.* 2003;146:351-358.
 72. **Brunner EJ, Shipley MJ, Witte DR, Fuller JH, Marmot MG.** Relation between blood glucose and coronary mortality over 33 years in the Whitehall Study. *Diabetes Care.* 2006;29:26-31.
 73. **Hu FB, Stamper MJ, Haffner SM, Solomon CG, Willett WC, Manson JE.** Elevated risk of cardiovascular disease prior to clinical diagnosis of type 2 diabetes. *Diabetes Care.* 2002;25:1129-1134.
 74. **Rijkkelijkhuizen JM, Nijpels G, Heine RJ, Bouter LM, Stehouwer CD, Dekker JM.** High risk of cardiovascular mortality in individuals with impaired fasting glucose is explained by conversion to diabetes: the Hoorn study. *Diabetes Care.* 2007;30:332-336.
 75. **Lorenzo C, Okoloise M, Williams K, Stern MP, Haffner SM.** San Antonio Heart Study. The metabolic syndrome as predictor of type 2 diabetes: the San Antonio heart study. *Diabetes Care.* 2003;26:3153-3159.
 76. **Ford ES, Zhao G, Li C.** Pre-diabetes and the risk for cardiovascular disease. *J Am Coll Cardiol.* 2010;55:1310-1317.
 77. **Pérez CM, Soto-Salgado M, Suárez E, Guzmán M, Ortiz AP.** High prevalence of diabetes and prediabetes and their coexistence with cardiovascular risk factors in a Hispanic community. *J Immigr Minor Health.* 2015;17:1002-1009.
 78. **Sörenson BM, Houben AJHN, Berendschot TTJM, et al.** Hyperglycemia is the main mediator of prediabetes- and type 2 diabetes-associated impairment of microvascular function: The Maastricht Study. *Diabetes Care.* 2017;40:e103-e105.
 79. **Motillo S, Filion KB, Genest J, et al.** The metabolic syndrome and cardiovascular risk. *J Am Coll Cardiol.* 2010;56:1110-1132.
 80. **Grundy SM.** Pre-diabetes, metabolic syndrome, and cardiovascular risk. *J Am Coll Cardiol.* 2012;59:635-643.
 81. **Vistisen D, Witte DR, Brunner EJ, et al.** Risk of cardiovascular disease and death in individuals with prediabetes defined by different criteria: the Whitehall II Study. *Diabetes Care.* 2018;41:899-906.
 82. **Ratner R, Goldberg R, Haffner S, et al.** Impact of intensive lifestyle and metformin therapy on cardiovascular disease risk factors

- in the diabetes prevention program. *Diabetes Care.* 2005;28:888-894.
83. **DECODE Study Group, European Diabetes Epidemiology Group.** Is the current definition for diabetes relevant to mortality risk from all causes and cardiovascular and noncardiovascular diseases? *Diabetes Care.* 2003;26:688-696.
 84. **Qiao Q, Jousilahti P, Eriksson J, Tuomilehto J.** Predictive properties of impaired glucose tolerance for cardiovascular risk are not explained by the development of overt diabetes during follow-up. *Diabetes Care.* 2003;26:2910-2914.
 85. **Rodriguez BL, Lau N, Burchfiel CM, et al.** Glucose intolerance and 23-year risk of coronary heart disease and total mortality: the Honolulu Heart Program. *Diabetes Care.* 1999;22:1262-1265.
 86. **Salimi Y, Fotouhi A, Mohammad K, Mansournia N, Mansournia MA.** Causal effects of intensive lifestyle and metformin interventions on cardiovascular disease risk factors in prediabetic people: an application of G-estimation. *Arch Iran Med.* 2017;20:55-59.
 87. **Siddiqui F, Kurbasic A, Lindblad U, Nilsson PM, Bennet L.** Effects of a culturally adapted lifestyle intervention on cardiometabolic outcomes: a randomized controlled trial in Iraqi immigrants to Sweden at high risk for type 2 diabetes. *Metabolism.* 2017;66:1-13.
 88. **Su W, Chen F, Dall TM, Iacobucci W, Perreault L.** Return on investment for digital behavioral counseling in patients with prediabetes and cardiovascular disease. *Prev Chronic Dis.* 2016;13:E13.
 89. **Nanditha A, Ram J, Snehalatha C, et al.** Early improvement predicts reduced risk of incident diabetes and improved cardiovascular risk in prediabetic Asian Indian men participating in a 2-year lifestyle intervention program. *Diabetes Care.* 2014;37:3009-3015.
 90. **Lindström J, Ilanne-Parikka P, Peltonen M, et al.** Finnish Diabetes Prevention Study Group. Sustained reduction in the incidence of type 2 diabetes by lifestyle intervention: follow-up of the Finnish Diabetes Prevention Study. *Lancet.* 2006;368:1673-1679.
 91. **Orchard TJ, Temprosa M, Goldberg R, et al.** Diabetes Prevention Program Research Group. The effect of metformin and intensive lifestyle intervention on the metabolic syndrome: the Diabetes Prevention Program randomized trial. *Ann Intern Med.* 2005;142:611-619.
 92. **Uusitupa M, Lindi V, Louheranta A, et al.** Finnish Diabetes Prevention Study Group. Long-term improvement in insulin sensitivity by changing lifestyles of people with impaired glucose tolerance: 4-year results from the Finnish Diabetes Prevention Study. *Diabetes.* 2003;52:2532-2538.
 93. **Ackermann RT, Marrero DG, Hicks KA, et al.** An evaluation of cost sharing to finance a diet and physical activity intervention to prevent diabetes. *Diabetes Care.* 2006;29:1237-1241.
 94. **Dalziel K, Segal L.** Time to give nutrition interventions a higher profile: a cost-effectiveness of 10 nutrition interventions. *Health Promot Int.* 2007;22:271-283.
 95. **Palmer AJ, Roze S, Valentine WJ, Spinaz GA, Shaw JE, Zimmet PZ.** Intensive lifestyle changes or metformin in patients with impaired glucose tolerance: modeling the long term health economic implications of the diabetes prevention program in Australia, France, Germany, Switzerland, and the United Kingdom. *Clin Ther.* 2004;26:304-321.
 96. **Lindgren P, Lindström J, Tuomilehto J, et al.** DPS Study Group. Lifestyle intervention to prevent diabetes in men and women with impaired glucose tolerance is cost-effective. *Int J Technol Assess Health Care.* 2007;23:177-183.
 97. **Ramachandran A, Snehalatha C, Yamuna A, Mary S, Ping Z.** Cost-effectiveness of the interventions in the primary prevention of diabetes among Asian Indians: within trial results of the Indian Diabetes Prevention Programme (IDPP). *Diabetes Care.* 2007;30:2548-2552.
 98. **Diabetes Prevention Program Research Group.** Within trial cost-effectiveness of lifestyle intervention or metformin for the primary prevention of type 2 diabetes. *Diabetes Care.* 2003;26:2518-2523.
 99. **Mudaliar U, Zabetian A, Goodman M, et al.** Cardiometabolic risk factor changes observed in diabetes prevention programs in US settings: a systematic review and meta-analysis. *PLoS Med.* 2016;13:e1002095.
 100. **Boyko EJ, Fujimoto WY, Leonetti DL, Newell-Morris L.** Visceral adiposity and risk of type 2 diabetes: a prospective study among Japanese Americans. *Diabetes Care.* 2000;23:465-471.
 101. **Neeland IJ, Turer AT, Ayers CR, et al.** Dysfunctional adiposity and risk of prediabetes and type 2 diabetes in obese adults. *JAMA.* 2012;308:1150-1159.
 102. **Gupta AK, Ravussin E, Johannsen DL, Stull AJ, Cefalu WT, Johnson WD.** Endothelial dysfunction: an early cardiovascular risk marker in asymptomatic obese individuals with prediabetes. *Br J Med Res.* 2012;2:413-423.
 103. **Weir GC, Bonner-Weir S.** Five stages of evolving beta-cell dysfunction during progression to diabetes. *Diabetes.* 2004;53(suppl 3):S16-S21.
 104. **Festa A, Williams K, D'Agostino R Jr, Wagenknecht LE, Haffner SM.** The natural course of beta-cell function in nondiabetic and diabetic individuals—the insulin resistance atherosclerosis study. *Diabetes.* 2006;55:1114-1120.
 105. **Gastaldelli A, Ferrannini E, Miyazaki Y, Matsuda M, DeFronzo RA.** Beta-cell dysfunction and glucose intolerance: results from the San Antonio metabolism (SAM) study. *Diabetologia.* 2004;47:31-39.
 106. **Butler AE, Janson J, Bonner-Weir S, Ritzel R, Rizza RA, Butler PC.** Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. *Diabetes.* 2003;52:102-110.
 107. **Kloppel G, Löhr M, Habich K, Oberholzer M, Heitz PU.** Islet pathology and the pathogenesis of type 1 and type 2 diabetes mellitus revisited. *Surv Synth Pathol Res.* 1985;4:110-125.
 108. **Ward WK, Bolgiano DC, McKnight B, Halter JB, Porte D Jr.** Diminished B cell secretory capacity in patients with noninsulin dependent diabetes mellitus. *J Clin Invest.* 1984;74:1318-1328.
 109. **Deng S, Vatamaniuk M, Huang X, et al.** Structural and functional abnormalities in the islets isolated from type 2 diabetic subjects. *Diabetes.* 2004;53:624-632.
 110. **Del Guerra S, Lupi R, Marselli L, et al.** Functional and molecular defects of pancreatic islets in human type 2 diabetes. *Diabetes.* 2005;54:727-735.
 111. **Gujral UP, Mohan V, Pradeepa R, et al.** Ethnic variations in diabetes and prediabetes prevalence and the roles of insulin resistance and b-cell function: the CARRS and NHANES studies. *J Clin Transl Endocrinol.* 2016;4:19-27.
 112. **Do OH, Gunton JE, Gaisano HY, Thorn P.** Changes in beta cell function occur in prediabetes and early disease in the Leprdb mouse model of diabetes. *Diabetologia.* 2016;59:1222-1230.
 113. **Malin SK, Solomon TPJ, Blaszczak A, Finnegan S, Filion J, Kirwan JP.** Pancreatic b-cell function increases in a linear dose-response manner following exercise training in adults with prediabetes. *Am J Physiol Endocrinol Metab.* 2013;305:E1248-E1254.
 114. **Kim SH, Liu A, Ariel D, et al.** Pancreatic beta cell function following liraglutide-augmented weight loss in individuals with prediabetes: analysis of a randomized placebo-controlled study. *Diabetologia.* 2014;57:455-462.
 115. **Ji Q.** Treatment strategy for type 2 diabetes with obesity: focus on glucagon-like peptide-1 receptor agonists. *Clin Ther.* 2017;39:1244-1264.
 116. **Garvey WT.** Comment on Cefau et al. update and next steps for real-world translation of interventions for type 2 diabetes prevention: reflections from a diabetes care editors' forum. *Diabetes Care.* 2016;39:1186-1201. *Diabetes Care.* 2017;40:e21-e22.
 117. **Li, M, Zeng T, Liu R, Chen L.** Detecting tissue-specific early warning signals for complex diseases based on dynamical network biomarkers: study of type 2 diabetes by cross-tissue analysis. *Brief Bioinform.* 2014;15:229-243.
 118. **Jin B, Liu R, Hao S, et al.** Defining and characterizing the critical transition state prior to the type 2 diabetes disease. *PLoS One.* 2017;12:e0180937.
 119. **Khera AV, Emdin CA, Drake I, et al.** Genetic risk, adherence to a healthy lifestyle, and coronary disease. *N Engl J Med.* 2016;375:2349-2358.
 120. **Tuomilehto J, Lindström J, Eriksson JG, et al.** Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med.* 2001;344:1343-1350.

121. Pan XR, Li GW, Hu YH, et al. Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance: the Da Qing IGT and Diabetes Study. *Diabetes Care.* 1997;20:537-544.
122. Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V. The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia.* 2006;49:289-297.
123. Ramachandran A, Snehalatha C, Mary S, et al. Pioglitazone does not enhance the effectiveness of lifestyle modification in preventing conversion of impaired glucose tolerance to diabetes in Asian Indians: results of the Indian Diabetes Prevention Programme-2 (IDPP-2). *Diabetologia.* 2009;52:1019-1026.
124. Ramachandran A, Arun N, Shetty AS, Snehalatha C. Efficacy of primary prevention interventions when fasting and postglucose dysglycemia coexist. *Diabetes Care.* 2010;33:2164-2168.
125. Flanu A, Bourse L, Naty N, et al. Long-term effectiveness of a lifestyle intervention for the primary prevention of type 2 diabetes in a low socio-economic community – an intervention follow-up study on Reunion Island. *PLoS One.* 2016;11:e0146095.
126. Davies MJ, Gray LJ, Troughton J, et al. A community based primary prevention programme for type 2 diabetes integrating identification and lifestyle intervention for prevention: the Let's Prevent Diabetes cluster randomized controlled trial. *Prev Med.* 2016;84:48-56.
127. Hu Z, Qin L, Xu H. One-year results of a synthetic intervention model for the primary prevention of T2D among elderly individuals with prediabetes in rural China. *Int J Environ Res Public Health.* 2017;14:E417.
128. Phillips LS, Weintraub WS, Ziemer DC, et al. All pre-diabetes is not the same: metabolic and vascular risks of impaired fasting glucose at 100 versus 110 mg/dl. *Diabetes Care.* 2006;29:1405-1407.
129. Zhang X, Gregg EW, Williamson DF, et al. A1C level and future risk of diabetes: a systematic review. *Diabetes Care.* 2010;33:1665-1673.
130. Cusick M, Meleth AD, Agrón E, et al. Associations of mortality and diabetes complications in patients with type 1 and type 2 diabetes: early treatment diabetic retinopathy study report no. 27. *Diabetes Care.* 2005;28:617-625.
131. Egede MB, Henriksen J-E, Durck TT, et al. Glucose effectiveness in nondiabetic relatives: dysglycemia and b-cell function at 10 years. *J Clin Endocrinol Metab.* 2014;99:1420-1424.
132. Laiteerapong N, Cifu AS. Screening for prediabetes and type 2 diabetes mellitus. *JAMA.* 2016;315:697-698.
133. Palmer ND, Stevens RD, Antinozzi PA, et al. Metabolomic profile associated with insulin resistance and conversion to diabetes in the insulin resistance atherosclerosis study. *J Clin Endocrinol Metab.* 2015;100:E463-E468.
134. Danker R, Chetrit A, Shanik MH, Raz I, Roth J. Basal state hyperinsulinemia in health normoglycemic adults heralds dysglycemia after more than two decades of follow up. *Diabetes Metab Res Rev.* 2012;28:618-624.
135. Walker RJ, Smalls BL, Campbell JA, Strom Williams JL, Egede LE. Impact of social determinants of health on outcomes for type 2 diabetes: a systematic review. *Endocrine.* 2014;47:29-48.
136. Tanrikulu MA, Agirbasli M, Berenson G. Primordial prevention of cardiometabolic risk in childhood. *Adv Exp Med Biol.* 2017;956:489-496.
137. Garvey WT, Arathuzik G. Lifestyle therapy for diabetes mellitus. In: Mechanick JI, Kushner RF, eds. *Lifestyle Medicine.* New York: Springer; 2016: 221-243.
138. Garvey WT, Arathuzik G, Miller GD, et al. Lifestyle therapy in the management of cardiometabolic risk: diabetes prevention, hypertension, and dyslipidemia. In: Mechanick JI, Kushner RF, eds. *Lifestyle Medicine.* New York: Springer; 2016: 245-267.
139. Poveda A, Chen Y, Brändström A, et al. The heritable basis of gene-environment interactions in cardiometabolic traits. *Diabetologia.* 2017;60:442-452.
140. Tuomilehto J, Lindström J, Eriksson JG, et al. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med.* 2001;344:1343-1350.
141. Samson SL, Garber AJ. Prevention of type 2 diabetes mellitus: potential of pharmacological agents. *Best Pract Res Clin Endocrinol Metab.* 2016;30:357-371.
142. Rubino F, Nathan DM, Eckel RH, et al. Metabolic surgery in the treatment algorithm for type 2 diabetes: a joint statement by international diabetes organizations. *Obes Surg.* 2017;27:2-21.
143. van der Leeuw J, Beulens JW, van Dieren S, et al. Novel biomarkers to improve the prediction of cardiovascular event risk in type 2 diabetes mellitus. *J Am Heart Assoc.* 2016;5:e003048.
144. Mechanick JI, Kushner RF. Why lifestyle medicine? In: Mechanick JI, Kushner RF, eds. *Lifestyle Medicine.* New York: Springer; 2016: 1-8.
145. Scales R, Buman MP. Paradigms of lifestyle medicine and wellness. In: Mechanick JI, Kushner RF, eds. *Lifestyle Medicine.* New York: Springer; 2016: 29-40.
146. Fletcher RH, Fletcher SW, Fletcher GS. *Clinical Epidemiology: The Essentials.* 5th ed. Philadelphia, PA: Wolters Kluwer/Lippincott Williams & Wilkins; 2014.
147. Hopman P, de Bruin SR, Forjaz MJ, et al. Effectiveness of comprehensive care programs for patients with multiple chronic conditions or frailty: a systematic literature review. *Health Policy.* 2016;120:818-832.
148. Knowles SE, Chew-Graham C, Adeyemi I, Coupe N, Coventry PA. Managing depression in people with multimorbidity: a qualitative evaluation of an integrated collaborative care model. *BMC Fam Pract.* 2015;16:32.
149. Langwell K, Keene C, Zullo M, Ogu LC. An American Indian community implements the chronic care model: evolution and lessons learned. *Health Promot Pract.* 2014;15(suppl 2):23S-28S.
150. Ku GMV, Kegels G. Implementing elements of a context-adapted chronic care model to improve first-line diabetes care: effects on assessment of chronic illness care and glycaemic control among people with diabetes enrolled to the First-Line Diabetes Care (FiLDCare) Project in the Northern Philippines. *Prim Health Care Res Dev.* 2015;16:481-491.
151. Admiraal WM, Holleman F, Snijder MB, et al. Ethnic disparities in the association of impaired fasting glucose with the 10-year cumulative incidence of type 2 diabetes. *Diab Res Clin Pract.* 2014;103:127-132.
152. Kim YA, Ku EJ, Khang AR, et al. Role of various indices derived from an oral glucose tolerance test in the prediction of conversion from prediabetes to type 2 diabetes. *Diabetes Res Clin Pract.* 2014;106:351-359.
153. Dauvrial M, Lorant V, d'Hoore W. Is the chronic care model integrated into research examining culturally competent interventions for ethnically diverse adults with type 2 diabetes mellitus? A review. *Eval Health Prof.* 2015;38:435-463.
154. Sepers CE, Fawcett SB, Lipman R, Schultz J, Colie-Akers V, Perez A. Measuring the implementation and effects of a coordinated care model featuring diabetes self-management education within four patient-centered medical homes. *Diab Educ.* 2015;41:328-342.
155. Eisenstat SA, Chang Y, Porneala BC, et al. Development and implementation of a collaborative team care model for effective insulin use in an academic medical center primary care network. *Am J Med Qual.* 2017;32:397-405.
156. Chung WW, Cua SS, Lai PSM, Chan SP. Effects of a pharmaceutical care model on medication adherence and glycaemic control of people with type 2 diabetes. *Patient Prefer Adherence.* 2014;8:1185-1194.
157. Baptista DR, Wiens A, Pontarolo R, Regis L, Reis WC, Correr CJ. The chronic care model for type 2 diabetes: a systematic review. *Diabetol Metab Syndr.* 2016;8:7.
158. Bodenheimer T, Willard-Grace R. The chronic care model and the transformation of primary care. In: Mechanick JI, Kushner RF, eds. *Lifestyle Medicine.* New York: Springer; 2016: 89-96.
159. Bloomgarden ZT, Einhorn DE. Hemoglobin A1c in diabetes diagnosis: time for caution. *Endocr Pract.* 2010;16:5-6.
160. Cobb J, Eckhart A, Perichon R, et al. A novel test for IGT utilizing metabolite markers of glucose tolerance. *J Diab Sci Technol.* 2015;9:69-76.

161. **Gerstein HC, Paré G, McQueen MJ, et al.** Identifying novel biomarkers for cardiovascular events or death in people with dysglycemia. *Circulation.* 2015;132:2297-2304.
162. **Millar SR, Perry IJ, Phillips CM.** HbA1c alone is a poor indicator of cardiometabolic risk in middle-aged subjects with pre-diabetes but is suitable for type 2 diabetes diagnosis: a cross-sectional study. *PLoS One.* 2015;10:0134154.
163. **de Graaf G, Postmus D, Bakker SJJ, Buskens E.** Design of stepwise screening for prediabetes and type 2 diabetes based on costs and cases detected. *J Clin Epidemiol.* 2015;68:1010-1018.
164. **McCoy RG, Van Houten HK, Ross JS, Montori VM, Shah ND.** HbA1c overtesting and overtreatment among US adults with controlled type 2 diabetes, 2001-13: observational population based study. *BMJ.* 2015;351:h6138.
165. **Guo F, Garvey WT.** Development of a weighted cardiometabolic disease staging (CMDS) system for the prediction of future diabetes. *J Clin Endocrinol Metab.* 2015;100:3871-3877.
166. **Guo F, Garvey WT.** Cardiometabolic disease staging predicts effectiveness of weight loss therapy to prevent type 2 diabetes: pooled results from phase 3 clinical trials assessing phentermine/topiramate-extended release. *Diabetes Care.* 2017, Apr 28 [Epub ahead of print].
167. **Desmedt M, Vertriest S, Hellings J, et al.** Economic impact of integrated care models for patients with chronic diseases: a systematic review. *Value Health.* 2016;19:892-902.
168. **Kirsch F.** Economic evaluations of multicomponent disease management programs with Markov models: a systematic review. *Value Health.* 2016;19:1039-1054.
169. **Seuring T, Archangelidi O, Suhrcke M.** The economic costs of type 2 diabetes: a global systematic review. *Pharmacoeconomics.* 2015;33:811-831.
170. **López-Martínez N, Brosa M, Segú-Tolsà L, Kalfhaus L, Bohigas L.** Efficiency of a personalized care model in diabetes as an example of chronic disease with information and communications technology support. *Value Health.* 2015;18:A617.
171. **McGinley EL, Gabbay RA.** The impact of new payment models on quality of diabetes care and outcomes. *Curr Diab Rep.* 2016;16:51.
172. **Van Hecke A, Heinen M, Fernández-Ortega P, et al.** Systematic literature review on effectiveness of self-management support interventions in patients with chronic conditions and low socioeconomic status. *J Adv Nurs.* 2017;73:775-793.
173. **Dagogo-Jack S.** 2015 Presidential Address: 75 years of battling diabetes – our global challenge. *Diabetes Care.* 2016;39:3-9.
174. **Schneider EC, Zaslavsky AM, Epstein AM.** Racial disparities in the quality of care for enrollees in Medicare managed care. *JAMA.* 2002;287:1288-1294.
175. **Mechanick JI, Harrell RM, Allende-Vigo MZ, et al.** Transculturalization recommendations for developing Latin American clinical practice algorithms in endocrinology – proceedings of the 2015 Pan-American workshop by the American Association of Clinical Endocrinologists and American College of Endocrinology. *Endocr Pract.* 2016;22:476-501.
176. **Mechanick JI, Leroith D.** Synthesis: deriving a core set of recommendations to optimize diabetes care on a global scale. *Ann Glob Health.* 2015;81:874-883.
177. **Knoops KT, de Groot LC, Kromhout D, et al.** Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women. *JAMA.* 2004;292:1433-1439.
178. **Estruch R, Ros E, Salas-Salvado J, et al.** Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med.* 2013;368:1279-1290.
179. **Kernan WN, Viscoli CM, Dearborn JL, et al.** Targeting pioglitazone hydrochloride therapy after stroke or transient ischemic attack according to pretreatment risk for stroke or myocardial infarction. *JAMA Neurol.* 2017;74:1319-1327.
180. **Gong Q, Zhang P, Wang J, et al.** Changes in mortality in people with IGT before and after the onset of diabetes during the 23-year follow-up of the Da Qing Diabetes Prevention Study. *Diabetes Care.* 2016;39:1550-1555.
181. **Vas PRJ, Alberti KG, Edmonds ME.** Prediabetes: moving away from a glucocentric definition. *Lancet Diab Endocrinol.* 2017;5:848-849.
182. **Justin Rucker A, Crowley SD.** The role of macrophages in hypertension and its complications. *Pflugers Arch.* 2017;469:419-430.